

SURVIVABILITY

newborn lambs and kids.

"Average lamb mortality rates have changed very little over the past few decades staying around 15%, of which 50% will occur in the first 24 hours after birth."

colostrum supplement or replacer for

Receiving a colostrum feeding as soon as possible after birth has been identified as being the single-most important step for newborn lamb and kid survival.

In addition, Colostrx LK can provide high immunity levels to newborn lambs and kids which optimize long-term performance, productivity and health under a variety of environmental or management conditions.

¹R. Wonfor, IBERS, Aberystwyth University, Improving lamb survival, Improving lamb survival



ENERGY

Newborn lambs and kids have limited energy reserves and need rapid access to colostrum to maintain body temperature and survive. This is especially true for lambs or kids that have a difficult birth, are born into cold temperatures, or are born to poorly fed ewes or does.

Colostrx LK contains natural high levels of colostral fat and lactose because they are the only nutrients that newborn lambs and kids can quickly convert into the energy needed to maintain body temperature.

We never strip out the natural colostral fat and replace it with animal fat, coconut oil, vegetable oil or other fat sources that offer minimal benefit to lambs and kids.





THE COLOSTRUM BRAND YOU CAN TRUST

We ensure safety, efficacy, purity and potency as a USDA Veterinary Biologics permitted product. We also rigorously test our colostrum powder to ensure the products are clean and effectively heat-treated to eradicate any contaminants from the sourced maternal colostrum.

EASY TO USE

Colostrx LK can be easily mixed in 15 seconds!

Check the back of the package for proper mixing instructions and remember to mix with water that is 110 to 120°F.

DIRECTIONS FOR USE

Mix the powder with clean, warm water 110 to 120°F [43 to 49°C]. Newborn lambs and kids should receive their first feeding of colostrum as soon after birth as possible by nipple bottle or esophageal tube.

Each feeding is based on body weight. Feed the appropriate amount described in the table 3 times in the first 18 hours of life.

Lamb or Kid weight / Peso del cordero o cabrito		Powder per feeding / Polvo por ración		Approx. # of scoops / Aprox. numero de cucharadas	Water per feeding / Agua por ración	
lb	kg	OZ	g	-	OZ	mL
5.00	2.30	0.90	25.00	1.25	1.60	48.00
6.00	2.70	1.10	30.00	1.50	1.90	57.00
7.00	3.20	1.20	35.00	1.75	2.30	67.00
8.00	3.60	1.40	40.00	2.00	2.60	76.00
9.00	4.10	1.60	45.00	2.25	2.90	86.00
10.00	4.60	1.80	50.00	2.50	3.20	95.00
11.00	5.00	1.90	55.00	2.75	3.60	105.00

60.00

3.00

COLOSTRX

BOVINE laG





>5.00